

Effect of pre-harvest fruit bagging on development and quality of mango cv Kesar under western Maharashtra conditions

SB JADHAV¹, VK GARANDE^{1*} and VK BHALERAO²

¹Regional Fruit Research Station, Ganeshkhind, Pune 411067 Maharashtra, India

²Regional Extension Centre, College of Agriculture, Pune 411005 Maharashtra, India

*Email for correspondence: vishnugarande@gmail.com

© Society for Advancement of Human and Nature (SADHNA)

Received: 27.09.2025/Accepted: 29.10.2025

ABSTRACT

The present investigations were carried out at the Department of Horticulture, Mahatma Phule Krishi Vidyapeeth, Rahuri, Maharashtra during 2019-2022 to study the effect of pre-harvest bagging on the quality of mango fruits (cv Kesar). The experiment was laid out in randomized block design with eight treatments and three replications, including different bag types viz newspaper bag, brown paper bag, brown paper bag with polythene coating, skirting bag, polythene bag, butter paper bag, muslin cloth bag and control (no bagging). The results revealed that bagging had a significant effect on the quality of mango fruits. Among all treatments, skirting bag recorded the highest fruit weight (226.86 g), TSS (20.49 °Brix), total sugars (13.93%) and reducing sugars (5.12%), along with the best overall acceptability (8.35). Brown paper bag and butter paper bag also performed better with higher fruit weight, improved biochemical parameters and extended shelf life. Bagging markedly reduced fruit fly infestation, which was highest in control (12.88%) and negligible in most other treatments. However, no significant effect of bagging was observed on stem-end rot incidence. Economically, the skirting bag treatment recorded the highest net return (Rs 161,384/ha) and benefit-cost ratio (2.70), followed by butter paper bag. The study concludes that pre-harvest fruit bagging, particularly with skirting and brown paper bags, enhances fruit quality, reduces pest infestation and improves profitability in mango cv Kesar under western Maharashtra conditions.

Keywords: Mango; pre-harvest bagging; fruit quality; benefit-cost ratio

INTRODUCTION

Mango (*Mangifera indica* L) is unarguably considered as 'King' of all fruits due to its nutritional richness, unique taste, pleasant aroma and its religious and medicinal importance. In India, the production of mango was 20,772.33 thousand MT from an area of 2,350.30 thousand hectares and in Maharashtra the production was 459.15 thousand MT from an area of 164.40 thousand hectares in 2021-22 (Anon 2023). The area under mango cultivation is increasing every year but safe and quality mango production is not increased (Islam et al 2019a).

The way we grow fruit is undergoing a vital transformation. There is a critical and growing need for new, alternative methods to produce high-quality, appealing fruit while still managing diseases and pests. This shift is driven by two main factors: increasing

consumer anxiety over the use of synthetic agricultural chemicals and a greater focus on environmental awareness. Consequently, there is a much stronger emphasis on reducing the use of pesticides to safeguard the well-being of farm workers, protect consumer health and ensure environmental protection (Sharma et al 2009).

An attractive, spotless and pest free fruits fetch premium rate to the growers (Ravuri et al 2024). Mangoes are a beloved, seasonal treat, providing huge nutritional and economic value across tropical and subtropical nations. However, the systems that produce this vital crop are increasingly threatened by global challenges, especially a changing climate. To understand and manage this threat, scientists are heavily focused on phenology: the study of recurring natural events in a plant's life like flowering and fruiting and how those events relate to their environment.

Because of this, worldwide phenological monitoring has become an immensely valuable tool for assessing the true effects of climate change (Halder et al 2024).

Pre-harvest fruit bagging is a powerful yet simple technique that shields fruit directly on the tree (Fan and Mattheis 1998). This practice provides a crucial physical layer of protection for sensitive crops, creating an ideal micro-environment that ensures farmers can consistently produce high-quality, unblemished fruit with attractive colour (Yang et al 2009). This shield is vital because, during the growth phase, fruit is constantly under attack from biological and environmental stressors such as pests, diseases, sunscald and cracking, which would otherwise cause a drastic reduction in commercial value (Prasad et al 2016a, 2016b). Due to these substantial benefits, bagging has been used extensively around the world on many fruit crops, including mango, banana, guava, grape, apple and litchi (Kumar et al 2021). The technique successfully produces more attractive fruit with fewer blemishes and significantly reduces post-harvest losses, as documented in crops like mango (Sarker et al 2009, Ding and Syakirah 2010, Rahman et al 2019), guava (de Araújo Neto et al 2020, Sharma et al 2020) and banana (Lima et al 2020). Furthermore, pre-harvest bagging proves effective by reducing the incidence of diseases, insect-pest damage, mechanical injury, sunburn, fruit cracking and the presence of agrochemical residues on the fruit (Sharma et al 2014, Jakhar and Pathak 2016, Islam et al 2019b, Akter et al 2020).

The present research work was undertaken to study the effect of pre-harvest bagging on quality

of mango fruit cv Kesar under western Maharashtra conditions.

MATERIAL and METHODS

The present research was conducted at the instructional-cum-research farm of the Department of Horticulture, Mahatma Phule Krishi Vidyapeeth, Rahuri, Maharashtra during 2019 to 2022. The experiment was conducted in randomized block design with eight treatments replicated thrice with a unit of 50 fruits per treatment per replication. The uniformly grown mango fruits from 10 years old mango trees were bagged (Plate 1) at egg stage ie 60 days from fruit set with different types of bags (25 cm × 20 cm) which constituted the various treatments viz T₁: Newspaper bag, T₂: Brown paper bag, T₃: Brown paper bag with polythene coating, T₄: Skirting bag, T₅: Polythene bag, T₆: Butter paper bag, T₇: Muslin cloth bag and T₈: Control (no bagging). Perforations of 2 per cent were made at the edges of all the bags for proper ventilation required during fruit development except for skirting and muslin cloth bags. In case of more than two fruits in a cluster, the smaller, underdeveloped, deformed and spotted fruits were removed and only one healthy fruit was bagged. While bagging, the brown paper, newspaper, butter paper and plastic bags were stapled properly, so that these do not fell down and to prevent the entry of insect pests. The skirting and muslin cloth bags were tied with the help of thread. Before bagging, two perforations (≤ 4 mm diameter holes) were made for proper ventilation at the bottom of the polythene and brown paper bags with polythene coating. White and brown paper bags were not perforated as these bags automatically allowed



Plate 1. Mango fruits bagged with different types of bagging material

aeration. The observations on physical parameters of the fruits such as fruit length, fruit diameter, fruit weight, pulp to stone ratio, overall acceptability, shelf life of fruits; chemical parameters like, TSS, acidity, reducing sugars, non-reducing sugars and total sugars and incidence of fruit fly and stem-end rot were recorded. Four fruits were randomly selected per treatment per replication to record the various observations.

Growth parameters: The length from stalk end to the apex of fruit and diameter was measured with the help of a digital Vernier caliper. Weight of fruit, pulp and stone were recorded by using electronic weighing balance. The pulp to stone ratio was calculated from pulp weight and stone weight.

Chemical parameters: TSS content was determined by hand refractometer (ATAGO Company Ltd, Japan); titratable acidity by titrating a known volume of homogenized sample against 0.1 N NaOH using phenolphthalein as indicator. The results were expressed in per cent anhydrous citric acid (Anon 2005). The reducing, non-reducing and total sugars were estimated as per Lane and Eynon (1923) method with modification suggested by Ranganna (1977).

Sensory evaluation: The ripe fruits were examined for their sensory qualities for assessing colour, flavour and texture by panel of five judges with nine-point Hedonic scale (Amerine et al 1965).

Shelf life of fruits: Mature fruits were harvested at 80-85 per cent maturity. Twenty harvested mature fruits of each treatment were ripened at ambient temperature by using plastic crates with perforations and traditional paddy straw as ripening material. At the bottom, 2.5 cm layer of paddy straw was made on which fruits were arranged. Simultaneously, two more layers were kept on the first layer. The end of shelf life was noted when the fruits were spoiled.

Fruit fly infestation: The fruit infestation was determined by counting the total number of fruits and number of infested fruits and converted into per cent infestation as per below:

$$\text{Fruit fly infestation (\%)} = \frac{\text{Number of infested fruits}}{\text{Total number of fruits}} \times 100$$

Stem-end rot incidence: Disease incidence was measured 10 days after harvest. Anthracnose lesions

(caused by *Colletotrichum* spp) on the side of the fruit and stem-end rot lesions (caused by *Diplodia* spp) at the stem end of the fruit were rated for disease incidence.

Benefit-cost ratio: To assess the profitability level of mango based on bagging technology, simple tabular form and benefit-cost ratio (BCR) were checked. BCR was estimated by the ratio of gross return to variable cost and gross return to total cost.

Statistical analysis: The statistical analysis was performed as per the ANOVA suggested by Panse and Sukhatme (1985). The mean and standard deviations were computed using statistical software of OPSTAT programme.

RESULTS and DISCUSSION

Growth, yield and quality parameters: The data regarding growth parameters of fruits are presented in Table 1. The pooled data reveal that the average fruit weight was recorded higher in treatments T₄ (Skirting bag) (226.86 g), T₂ (Brown paper bag) (214.19 g) and T₆ (Butter paper bag) (212.40 g) which were at par as compared to lower weight observed in T₈ (Control) (180.44 g), T₅ (Polythene bag) (190.72 g), T₃: Brown paper bag with polythene coating (194.58 g), T₁ (Newspaper bag) (198.59 g) and T₇ (Muslin cloth bag) (199.94 g), all being at par. There was no effect of bagging on fruit length, fruit diameter and fruit yield per hectare.

Large number of physiological, biochemical and structural changes occur during the ripening of fruit which include the degradation of starch or other stored polysaccharides, production of sugars, synthesis of pigments and volatile compounds and the partial solubilization of cell wall (Jain et al 2003).

The pooled data regarding TSS content of the fruits presented in Table 2 show that higher TSS content was noticed in the treatment T₄ (20.49 °Brix), T₁ (19.95 °Brix), T₂ (19.61 °Brix) and T₅ (19.34 °Brix), which were at par, as compared to T₈ (17.84 °Brix), T₃ (18.41 °Brix), T₇ (18.89 °Brix) and T₆ (18.95 °Brix), which were also at par.

Total sugars were found to be higher in the fruits packed in T₄ (13.93%), T₂ (13.37%) and T₁ (13.30%), the three being at par, as compared to T₈ (12.11%), T₃ (12.56%), T₅ (12.61%) and T₇ (12.62%),

Table 1. Effect of bagging on growth and yield of mango fruits

| Treatment | Fruit length (cm) | Fruit diameter (cm) | Average fruit weight (g) | Fruit yield (tonnes/ha) |
|---|-------------------|---------------------|--------------------------|-------------------------|
| T ₁ : Newspaper bag | 10.94 | 6.98 | 198.59 | 6.76 |
| T ₂ : Brown paper bag | 11.08 | 6.86 | 214.19 | 7.29 |
| T ₃ : Brown paper bag with polythene coating | 10.39 | 6.70 | 194.58 | 7.04 |
| T ₄ : Skirting bag | 10.95 | 7.12 | 226.86 | 7.09 |
| T ₅ : Polythene bag | 10.47 | 6.76 | 190.72 | 6.56 |
| T ₆ : Butter paper bag | 10.47 | 6.85 | 212.40 | 7.18 |
| T ₇ : Muslin cloth bag | 10.47 | 6.85 | 199.94 | 6.48 |
| T ₈ : Control (no bagging) | 10.22 | 6.66 | 180.44 | 6.34 |
| SEm(±) | 0.20 | 0.12 | 7.87 | 0.40 |
| CD _{0.05} | NS | NS | 24.09 | NS |

NS = Non-significant

Table 2. Effect of bagging on quality of mango fruits

| Treatment | TSS (°Brix) | Acidity (%) | Total sugars (%) | Reducing sugars (%) | Non-reducing sugars (%) | Pulp (%) | Pulp to stone ratio | Shelf life of fruits (days) |
|---|-------------|-------------|------------------|---------------------|-------------------------|----------|---------------------|-----------------------------|
| T ₁ : Newspaper bag | 19.95 | 0.41 | 13.30 | 4.91 | 8.61 | 61.72 | 3.72 | 14.41 |
| T ₂ : Brown paper bag | 19.61 | 0.42 | 13.37 | 4.92 | 8.27 | 61.56 | 3.72 | 12.80 |
| T ₃ : Brown paper bag with polythene coating | 18.41 | 0.42 | 12.56 | 5.05 | 7.53 | 61.53 | 3.66 | 14.13 |
| T ₄ : Skirting bag | 20.49 | 0.42 | 13.93 | 5.12 | 8.47 | 61.97 | 3.67 | 14.39 |
| T ₅ : Polythene bag | 19.34 | 0.47 | 12.61 | 4.73 | 7.21 | 60.17 | 3.62 | 13.21 |
| T ₆ : Butter paper bag | 18.95 | 0.46 | 13.08 | 4.85 | 7.64 | 60.95 | 3.69 | 14.20 |
| T ₇ : Muslin cloth bag | 18.89 | 0.44 | 12.62 | 4.69 | 7.80 | 59.51 | 3.60 | 13.01 |
| T ₈ : Control (no bagging) | 17.84 | 0.49 | 12.11 | 4.57 | 7.60 | 58.25 | 3.46 | 12.19 |
| SEm(±) | 0.41 | 0.02 | 0.23 | 0.09 | 0.35 | 1.34 | 0.07 | 0.64 |
| CD _{0.05} | 1.26 | NS | 0.70 | 0.28 | NS | NS | NS | NS |

NS = Non-significant

which were also at par. Higher reducing sugars were observed in T₄ (5.12%), T₃ (5.05%), T₂ (4.92%), T₁ (4.91%) and T₆ (4.85%), all being at par and lower in T₈ (4.57%), T₇ (4.69%) and T₅ (4.73%) which were at par and also at par with T₆ (4.85%). There was no effect of bagging on the acidity, non-reducing sugars, pulp percentage, pulp to stone ratio and self life of fruits.

Bagging significantly influences the physical, biochemical and shelf life characteristics of mango fruits. Islam et al (2019b) reported that mango cv Langra fruits bagged with white and brown paper bags recorded the highest fruit length (97.93 and 103.5 mm), diameter (79.27 and 84.85 mm), weight (311.66 and 329.2 g) and pulp weight (278.9 and 289.8 g)

respectively, while the lowest values were observed in polythene bag and control. Afsar and Sultana (2019) also found brown paper bags most effective for fruit weight, length and diameter. Haldankar et al (2015) observed that newspaper and brown paper bags improved fruit weight, diameter and pulp weight in Alphonso mango. Kumari (2023) reported that brown paper bagging in Mallika enhanced fruit weight (630.08 g), fruit length (16.34 cm), diameter (8.35 cm) and volume (591.83 ml) while in Pusa Shrestha, white paper bagging improved fruit weight (358.37 g), length (12.39 cm), diameter (6.65 cm) and volume (327.83 ml). Veeramanikandan et al (2024) found that Banganapalli mangoes bagged in brown/black double-layered bags showed the highest fruit weight (641.8 g) and improved

peel colour and firmness. Ramez et al (2024) reported the highest fruit weight (256.00 g), volume (224.38 ml) and length (89.68 mm) in non-woven green bag treatment though chemical traits were not significantly affected.

Islam et al (2019b) and Haldankar et al (2015) observed higher total soluble solids (TSS), reducing and non-reducing sugars in paper-bagged fruits compared to control. Kireeti et al (2016) found that muslin cloth bags gave the highest TSS (17.61 °B), butter paper bags had maximum reducing (2.82%) and total sugars (9.65%) while polythene bags showed minimum values.

Sharma et al (2023) demonstrated that Dasherri fruits bagged in brown paper and treated with 0.05 per cent carbendazim had the highest TSS (21.16 °Brix) and sugar contents with the lowest acidity (0.11 per cent). Kumari (2023) reported higher TSS (22.67 °Brix), total sugars (14.64%) and non-reducing sugars (10.05%) in Mallika under brown paper bagging while white paper bagging improved biochemical traits in Pusa Shrestha. Veeramanikandan et al (2024) also recorded improved TSS (17.72 °Brix) and TSS-acid ratio (60.17) in double-layered bag treatments.

Bagging consistently extended fruit shelf life across studies. Islam et al (2019b) found that brown paper bags extended Langra mango shelf life to 11 days compared to 6 days in control. Hossain et al (2020)

reported that brown paper bags prolonged Amrapali mango shelf life to 15 days while un-bagged fruits lasted only 8 days. Kumari (2023) observed shelf life of 11.67 days in Mallika and 14.67 days in Pusa Shrestha under paper bagging. Kireeti et al (2016) found that fruits bagged with newspaper and brown paper bags had the longest shelf life (17 days) and Veeramanikandan et al (2024) reported a 16-day shelf life in double-layered brown/black bagged Banganapalli mangoes.

Overall acceptability: The data presented in Fig 1 indicate that the mean organoleptic score for acceptability was the highest in T₄ (8.35) followed by T₃ (8.00) and T₂ (7.90) and the lowest in T₈ (6.50) followed by T₇ (6.70) and T₆ (7.20).

Pest and disease incidence: The data presented in Table 3 reveal that, significantly higher infestation of fruit fly was recorded in T₈ (12.88%) as compared to T₁ (3.89%). However, T₈ (12.88%) was also at par with T₂ (7.55%), which on the other hand was also at par with T₁ (3.89%). The fruit fly infestation was not noticed in rest of the treatments. There was no effect of bagging on stem-end rot disease of mango.

Economics of bagging: The data regarding the economics of mango fruits bagging have been presented in Table 4. Highest net return (Rs 161,384) and B-C ratio (2.70) were recorded in T₄ followed by T₆ (Rs




Fig 1. Effect of bagging on overall acceptability score of mango fruits

147,704 and 2.25 respectively), whereas, the lowest in T_7 (Rs 57,761 and 1.25 respectively).

Bagging of mango fruits significantly influences sensory quality, pest and disease incidence and economic returns. Islam et al (2019b) reported improved sensory qualities in mango fruits bagged with brown and white paper bags compared to control. However, Haldankar et al (2015) observed no major change in sensory qualities of Alphonso fruits, though brown paper bags with polythene coating and butter paper, muslin cloth and scurting bags reduced the incidence of mealy bag. Kireeti et al (2016) ranked muslin cloth bags highest for sensory evaluation, while newspaper and brown paper bags provided the best shelf life (17 days). Akhilesh et al (2024) found non-woven and brown paper bags most effective for sensory attributes in Banganapalli mango, with the

highest scores for flavour (8.33), texture (8.67), taste (8.67) and overall acceptability (8.33).

Bagging also reduced pest and disease incidence in mango fruits. Hofman et al (1997) reported that bagging of cv Keitt decreased anthracnose and stem-end rot caused by *Colletotrichum* and *Dothoriella* spp. Veeramanikandan et al (2024) observed reduced fruit fly infestation and disease incidence (4.89%) in brown/black double-layered bags. del Pino et al (2021) found that yellow satin paper and white muslin cloth bags effectively minimized *Aulacaspis tubercularis* damage when applied before pest migration. Kumari (2023) reported that brown paper bagging in Mallika and white paper bagging in Pusa Shrestha minimized anthracnose, stem-end rot and sooty mould with lower incidence, severity and AUDPC values.

Table 3. Effect of fruit bagging on pest and disease incidence in mango

| Treatment | Disease incidence (%) | |
|--|-----------------------|---------------|
| | Fruit fly | Stem-end rot |
| T_1 : Newspaper bag | 3.89 (6.66) | 1.00 (3.32) |
| T_2 : Brown paper bag | 7.55 (13.06) | 3.33 (6.14) |
| T_3 : Brown paper bag with polythene coating | 0.00 | 0.00 |
| T_4 : Skirting bag | 0.00 | 0.00 |
| T_5 : Polythene bag | 0.00 | 6.66 (12.23) |
| T_6 : Butter paper bag | 0.00 | 0.00 |
| T_7 : Muslin cloth bag | 0.00 | 0.00 |
| T_8 : Control (no bagging) | 12.88 (21.0) | 10.11 (15.27) |
| S $\bar{E}m(+)$ | 3.56 | 4.35 |
| CD _{0.05} | 10.89 | NS |

NS = Non-significant; Figures in parentheses are angular transformations

Table 4. Economics of pre-harvest bagging materials

| Treatment | Pooled yield | Gross return | Cost of cultivation | Additional cost (Rs) | Total cost (Rs) | Net return (Rs) | BCR |
|--|--------------|--------------|---------------------|----------------------|-----------------|-----------------|------|
| T_1 : Newspaper bag | 6.76 | 304,200 | 145,948 | 16,427 | 162,375 | 141,825 | 1.87 |
| T_2 : Brown paper bag | 7.29 | 328,050 | 145,948 | 43,943 | 189,891 | 138,160 | 1.73 |
| T_3 : Brown paper bag with polythene coating | 7.04 | 352,000 | 145,948 | 31,958 | 177,906 | 138,894 | 1.98 |
| T_4 : Skirting bag | 7.09 | 425,400 | 145,948 | 11,718 | 157,666 | 161,384 | 2.70 |
| T_5 : Polythene bag | 6.56 | 295,200 | 145,948 | 26,632 | 172,580 | 122,620 | 1.71 |
| T_6 : Butter paper bag | 7.18 | 394,900 | 145,948 | 29,448 | 175,396 | 147,704 | 2.25 |
| T_7 : Muslin cloth bag | 6.48 | 291,600 | 145,948 | 87,885 | 233,833 | 57,767 | 1.25 |
| T_8 : Control (no bagging) | 6.34 | 285,300 | 145,948 | - | 145,948 | 139,352 | 1.95 |

Prices considered according to quality of fruits

Economically, bagging enhanced productivity and profitability. Afsar and Sultana (2019) recorded higher yield (10,850 kg/ha), gross return (Tk 596,750) and benefit-cost ratio (3.59) for adopters. Akhilesh et al (2024) observed increased yield (13,470 kg/ha) and the highest benefit-cost ratio (4.94) in reddish brown paper bag, while Kumari (2023) reported higher yield in Mallika (36.75 kg/tree, 14.70 t/ha) and Pusa Shrestha (21.14 kg/tree, 8.46 t/ha) with paper bagging.

CONCLUSION

Pre-harvest bagging significantly influenced the physical, biochemical and sensory quality of mango cv Kesar fruits. The use of skirting and brown paper bags resulted in superior fruit weight, higher TSS, sugars and overall acceptability, while effectively minimizing fruit fly infestation. Although no significant effect on stem-end rot was observed, bagging treatments enhanced fruit appearance and reduced blemishes. The skirting bag recorded the highest economic return and benefit-cost ratio, confirming its practical feasibility for commercial adoption. Thus pre-harvest fruit bagging, particularly with skirting or brown paper bags, can be recommended as a safe, eco-friendly and economically viable practice for improving marketable quality and profitability of mango production under western Maharashtra conditions.

REFERENCES

- Afsar M and Sultana N 2019. Effects of different bagging materials on mango production: assessment of profitability and fruit quality. *Journal of Agricultural and Rural Research* **4(1)**: 1-11.
- Akhilesh M, Harikanth P, Rajashekar M and Madhavi B 2024. Effect of pre-harvest fruit bagging on sensory and economics of mango cv Banganapalli. *International Journal of Research in Agronomy* **7(10S)**: 465-468.
- Akter MM, Islam MT, Akter N, Amin MF, Bari MA and Uddin MS 2020. Pre-harvest fruit bagging enhanced quality and shelf life of mango (*Mangifera indica* L) cv Amrapali. *Asian Journal of Agricultural and Horticultural Research* **5(3)**: 45-54.
- Amerine MA, Pangborn RM and Roessler EB 1965. Principles of sensory evaluation of food. In: *Food science and technology: a series of monographs* (ML Anson, CO Chichester, EM Mrak and GF Stewart, Eds), Academic Press, New York.
- Anonymous 2005. Official methods of analysis of AOAC International. 18th Edn, Association of Officiating Analytical Chemists, Washington, DC, USA.
- Anonymous 2023. Agricultural statistics at a glance 2023. Economics, Statistics and Evaluation Division, Department of Agriculture and Farmers' Welfare, Ministry of Agriculture and Farmers' Welfare, Government of India.
- de Araújo Neto SE, Rocha C, de Farias JF, Minosso SCC and Ferreira RLF 2020. Quality of guava fruits bagged with different materials in an organic system. *Comunicata Scientiae* **11**: e3206; doi: 10.14295/cs.v11i0.3206.
- del Pino M, Bienvenido C, Wong ME, del Carmen Rodríguez M, Boyero JR and Vela JM 2021. Influence of pre-harvest bagging on the incidence of *Aulacaspis tubercularis* Newstead (Hemiptera: Diaspididae) and fruit quality in mango. *Insects* **12**: 500; doi: 10.3390/insects12060500.
- Ding P and Syakirah MN 2010. Influence of fruit bagging on post-harvest quality of Harumanis mango (*Mangifera indica* L). *Acta Horticulturae* **877**: 169-174.
- Fan X and Mattheis JP 1998. Bagging 'Fuji' apples during fruit development affects colour development and storage quality. *HortScience* **33**: 1235-1238.
- Haldankar PM, Parulekar YR, Kireeti A, Kad MS, Shinde SM and Lawande KE 2015. Studies on influence of bagging of fruits at marble stage on quality of mango cv Alphonso. *Journal of Plant Studies* **4(2)**: 12-20.
- Halder R, Varma S, Singh M and Dahiya 2024. From orchard to table: understanding climate change impacts on mango production in India: a review. *International Journal of Advanced Biochemistry Research* **8(4)**: 472-478.
- Hofman PJ, Smith LG, Joyce DC, Johnson GI and Meiburg GF 1997. Bagging of mango (*Mangifera indica* cv Keitt) fruit influences fruit quality and mineral composition. *Post-harvest Biology and Technology* **12(1)**: 83-91.
- Hossain MJ, Hossain MM, Rabbani MG, Hafiz MMH and Islam MZ 2020. Effects of pre-harvest fruit bagging on post-harvest quality and shelf life of mango cv Amrapali. *Journal of Bangladesh Agricultural University* **18(1)**: 61-67.
- Islam MT, Shamsuzzoha M, Bari MA, Rahman MS, Akter MM, Islam M and Rahman MA 2019a. Influence of bagging time on fruit quality and shelf life of mango (*Mangifera indica* L) cv Amrapali in Bangladesh. *International Journal of Agriculture and Environmental Research* **5(4)**: 412-423.
- Islam MT, Shamsuzzoha M, Rahman MS, Bari MA, Akter MM, Khatun A, Huque R and Uddin MS 2019b. Effect of bagging time on fruit quality and shelf life of mango cv Langra in Bangladesh. *International Journal of Agriculture, Environment and Bioresearch* **4(4)**: 279-289.

- Jain N, Dhawan K, Malhotra S and Singh R 2003. Biochemistry of fruit ripening of guava (*Psidium guajava* L): compositional and enzymatic changes. *Plant Foods for Human Nutrition* **58**: 309-315.
- Jakhar MS and Pathak S 2016. Effect of pre-harvest nutrients application and bagging on quality and shelf life of mango (*Mangifera indica* L) fruits cv Amrapali. *Journal of Agricultural Science and Technology* **18(3)**: 717-729.
- Kireeti A, Haldankar PM, Babu MRV and Parulekar YR 2016. Effect of pre-harvest bagging on mango fruit quality. *Research in Environment and Life Sciences* **9(11)**: 1366-1369.
- Kumar M, Singh VP, Jat R, Ahamad S and Kumar V 2021. Pre-harvest fruit bagging for quality improvement in fruit crops: a review. *Pharma Innovation* **10(7)**: 530-541.
- Kumari I 2023. Effect of pre-harvest bagging on yield, quality and shelf life of mango (*Mangifera indica* L) cvs Mallika and Pusa Shrestha. MSc Thesis, GB Pant University of Agriculture and Technology, Pantnagar, Uttarakhand, India.
- Lane JH and Eynon L 1923. Determination of reducing sugar by means of Fehling's solution with methylene blue as internal indicator. *Journal of the Society of Chemical Industry* **17**: 32-37.
- Lima JD, Engelking EW, Rozane DE, Gomes EN, da Silva SHMG and Kluge RA 2020. Effect of bunch protection material and bagging time on the yield of Nanica banana and chilling control. *Australian Journal of Crop Science* **14(4)**: 574-580.
- Panse VG and Sukhatme PV 1985. *Statistical methods for agricultural workers*. 4th Edn, Indian Council of Agricultural Research, New Delhi, India, 359p.
- Prasad K, Sharma RR and Srivastav M 2016a. Post-harvest treatment of antioxidant reduces lenticels browning and improves cosmetic appeal of mango (*Mangifera indica* L) fruits without impairing quality. *Journal of Food Science and Technology* **53(7)**: 2995-3001.
- Prasad K, Sharma RR, Srivastav M and Sethi S 2016b. Effect of hot water treatment on the incidence of lenticels browning and quality of mango fruits. *Indian Journal of Horticulture*, **73(4)**: 576-581.
- Rahman MA, Alam SMK, Reza MH, Uddin MS, Amin MN and Nasrin TAA 2019. Impact of pre-harvest fruit bagging and improved post-harvest practices in reducing losses and managing quality of mango in the value chain system. *International Journal of Post-harvest Technology and Innovation* **6(2)**: 117-136.
- Ramez N, Subba S, Taheri MN, Alam M and Hasan MA 2024. Studies on the effect of bagging on yield and quality of mango cv Himsagar. *International Journal of Economic Plants* **11(4)**: 521-524.
- Ranganna S 1977. *Manual of analysis of fruit and vegetable products*. Tata McGraw-Hill, New York, USA, 634p.
- Ravuri PL, Deva S and Jyosthna MK 2024. Assessment of pre-harvest fruit bagging to manage pests of mango (*Mangifera indica* L) cv Benishan. *Asian Journal of Microbiology, Biotechnology and Environmental Sciences* **26(2)**: 211-214.
- Sarker D, Rahman MM and Barman JC 2009. Efficacy of different bagging materials for the control of mango fruit fly. *Bangladesh Journal of Agricultural Research* **34(1)**: 165-168.
- Sharma RR, Nagaraja A, Goswami AK, Thakre M, Kumar R and Varghese E 2020. Influence of on-the-tree fruit bagging on biotic stresses and post-harvest quality of rainy-season crop of Allahabad Safeda guava (*Psidium guajava* L). *Crop Protection* **135**: 105216; doi: 10.1016/j.cropro.2020.105216.
- Sharma RR, Reddy SVR and Jhalegar MJ 2014. Pre-harvest fruit bagging: a useful approach for plant protection and improved post-harvest fruit quality – a review. *Journal of Horticultural Science and Biotechnology* **89(2)**: 101-113.
- Sharma RR, Singh D and Singh R 2009. Biological control of post-harvest diseases of fruits and vegetables by microbial antagonists. *Biological Control* **50(3)**: 205-221.
- Sharma S, Malik S, Kumar R, Ujjwal V, Chand G, Sinha BK and Kumar R 2023. Effect of pre-harvest bagging and fungicide application on mango (*Mangifera indica* L) chemical characteristics. *Pharma Innovation* **12(11)**: 1607-1613.
- Veeramanikandan M, Muthuvel I, Soman P, Shanmugasundaram KA, Senthil A and Suganthi A 2024. Pre-harvest bagging: optimizing physical and quality attributes of Banganapalli mango (*Mangifera indica* L). *Plant Science Today* **11(SP4)**: doi: 10.14719/pst.5302.
- Yang W-H, Zhu X-C, Bu J-H, Hu G-B, Wang H-C and Huang X-M 2009. Effects of bagging on fruit development and quality in cross-winter off-season longan. *Scientia Horticulturae* **120**: 194-200.